


CARBON FOR CROPS

Assessing Impacts of Regenerative Organic Agriculture

Tyler Cobian, Ilayda Dinc, Alex Ivina, Karan Shetty
 Faculty advisor: Kyle Meng




BACKGROUND

Conventional agriculture is a massive contributor to greenhouse gas emissions, declines in soil health, and erosion of arable soil. These impacts will only worsen as the global population grows and the effects of climate change becomes more severe.

Regenerative organic agriculture is proposed as an alternative to conventional agriculture, with practices that aim **to improve soil health and reduce greenhouse gas emissions by allowing crops to sequester more carbon in the soil.** Some organizations, such as Patagonia Inc., aim to pair regenerative practices with existing organic standards. This type of system would use organic standards to keep synthetic inputs and GMOs out of the soil while promoting soil health with regenerative practices.

Patagonia is interested in pursuing regenerative organic agriculture in the production of crops they purchase. They hope to reduce their carbon footprint through regenerative agriculture, and want to quantify the carbon sequestration potential of regenerative organic agriculture in their supply chain. The analysis was conducted for the following crops in Patagonia's fiber and food production:



OBJECTIVES

- 1 Model soil carbon sequestration and GHG emissions of four crops under organic and regenerative management over multiple global locations.
- 2 Determine which regenerative practices had the greatest impact on soil carbon sequestration and greenhouse gas emissions.
- 3 Recommend to Patagonia which practices have the largest carbon sequestration and the lowest greenhouse gas emissions.

APPROACH

The team used the University of New Hampshire's DNDC soil model to simulate soil carbon and GHG emissions between organic and regenerative management for the four crops. This model required data inputs on:


- Climate:** For domestic sites, this data came from NOAA, while data for international sites was obtained from online resources.
- Soil properties:** This data for all crops and locations assessed came from ISRIC's SoilGrids, a meta analysis on global soil properties and carbon stock.
- Crop management practices:** This involved data on crop type, crop rotations practiced, tillage use, fertilizer use, irrigation and flooding, and the use of grazing.

Using this data, the DNDC model simulated soil carbon and nitrogen dynamics, as well as the emissions of trace gases, for each crop and location under a **20 year time period.**



RESULTS

Soil Organic Carbon




Total changes in soil organic carbon under regenerative and organic management after 20 years. **It can be seen that for all crops analyzed regenerative practices lead to more carbon stored in the soil after 20 years.**

All perennial crops (i.e. mango trees, grasslands, Kernza wheat) sequestered carbon. The annual crop (cotton) lost carbon, although it lost less carbon under regenerative management.


When total GHG emissions are taken into account, emissions are reduced in all crops except for grazing on perennial grasslands.

Some regenerative practices (such as compost use) can sequester carbon, but emit other greenhouse gases like nitrous oxide.

Net GHG Emissions



Location and GHG Emissions



Greenhouse gas emissions in cotton production in four locations. Regenerative management had lower emissions in all locations except China.

Regeneratives will not always produce lower emissions than organic. Practices will behave differently depending on climate and soil type.

KEY FINDINGS

- Regenerative organic agriculture builds soil organic carbon.
- Effects of regenerative agriculture on net greenhouse gas emissions are variable.
- The same practices will behave differently with variability in outcomes when implemented in different climate and soil properties.

RECOMMENDATIONS

Based on our results, we recommend Patagonia to **utilize perennial cover cropping and fiber crops**, as they grow deep root systems that allow them to sequester more carbon and maintain soil structure over time. Unlike annual crops, perennials do not need frequent replanting, which avoids both soil disturbance and carbon release.

Additionally, Patagonia should **pair regenerative practices that optimize carbon sequestration and GHG reduction**, such as combining compost/manure application and drip irrigation. This union would take advantage of the carbon benefits of compost while limiting the nitrous oxide emissions released from compost on over-watered fields.

To further understand the impact of regenerative organic agriculture, **yields should be closely monitored.** While no drop in yields was detected, such a decrease could increase greenhouse gas emissions and promote land clearing for additional cropping area.

ACKNOWLEDGEMENTS

We would like to thank our advisors Dr. Kyle Meng, Dr. Arturo Keller and Dr. Sangwon Suh as well as the Patagonia team Elissa Foster, Elena Egorova, and Stephanie Karba.

